

DEPRESSION AND CHILD NEGLECT IN MOTHERS OF CHILDREN WITH GROWTH RETARDATION

Idris YILDIRIM ¹, Hakan DEMIRCI ¹, Kaan DEMIROREN ² Emre YALCINTAS ¹

¹University of Health Sciences, Department of Family Medicine, Bursa Yuksek Ihtisas Training and Research Hospital, Bursa, Turkey.

²University of Health Sciences, Department of Pediatric, Bursa Yuksek Ihtisas Training and Research Hospital, Bursa, Turkey.

119

ABSTRACT

Aim: In this study, it was aimed to evaluate depression and child neglect in mothers of children with growth retardation (GR).

Methods: The research was carried out at Baęlaraltı Training Health Care Center affiliated to Bursa University of Health Sciences Yüksek Ihtisas Training and Research Hospital. The study is a case control study. 200 children with GR between the ages of 0-6 and their mothers' were enrolled as case group, as well as 200 children with similar age, gender, without GR and their mother were included in the study, as control group (CG). The Beck Depression Inventory (BDI) and the Maternal Child Neglect Scale were applied to the mothers.

Results: Depression levels of mothers whose children were in the GR group were high. BDI values increased the risk of GR by 1.062 times. In addition, it was found that physical neglect values increased the risk of GR by 1.464 times at a statistically significant level. The number of individuals living at home decreased the risk of GR and emotional neglect scores of the GR group.

Conclusions: In the study, the risk of depression and child neglect is higher in mothers of children with growth and development retardation. We believe that evaluating the mothers of children with developmental disabilities about depression and child neglect and taking the necessary measures can have a positive impact on the treatment processes of children.

Keywords: child neglect, emotional neglect, growth and development retardation, maternal depression, physical neglect

Corresponding Author: Idris Yildirim idris.yildirim@saglik.gov.tr

Received: December 31, 2024; **Accepted:** December 31, 2024; **Published Online:** December 31, 2024

Cite this article as: Yildirim, I., Demirci, H., Demiroren, K. & Yalcintas, E. (2024). Depression and Child Neglect in Mothers of Children With Growth Retardation. European Journal of Human Health 4(4),119-127.



INTRODUCTION

Currently, the survival of children during a healthy growth and development period depends on the interaction of many factors at different rates, such as genetic factors, as well as family relationships, the socioeconomic status of the family and society, nutritional conditions, and the mental health status of mothers. These various factors include important elements that determine the basic level of health and quality of life in the first years of children's life. Growth and development retardation may occur in many children, often due to many factors such as economic difficulties, malnutrition, domestic violence and neglect [1].

Growth retardation by definition; refers to the fact that the expected growth curves of children according to their age and gender remain below the standards of society. This condition occurs as a result of the interaction of genetic, psychosocial and environmental factors to different degrees. Many factors, such as nutritional deficiencies, genetic predisposition, recurrent infections and socioeconomic difficulties, can prevent children from realizing their full biological potential [2].

Maternal depression by definition; It is the appearance of signs of depression during the gestational period of mothers, puerperal pregnancy or early in the child's life. This situation negatively affects the mother's communication with her child and her maternal skills. Depression occurring in the mother can disrupt the dynamics within the family and

increase family tension, which negatively affects the healthy growth and development environment of the child, and as a result, retardation in the child's growth and development can be monitored [3]. Depression can reduce the mother's physical and emotional connection with her child, depending on this situation, the child may become vulnerable to child neglect. As a result of child neglect, it can have long-term consequences that can lead to biological, physical and psychosocial problems in the later years of a child's life [4].

This study aims to examine the complex relationships in this regard in more depth, focusing on understanding the interaction between depression and child neglect, especially in the mothers of children with growth and development Deceleration. This complexity represents an important area of research that can have long-term effects on child health and parent-child relationships [5].

METHODS

The research was carried out at Baglalaralti Training Health Care Center affiliated to Bursa University of Health Sciences Yuksek Ihtisas Training and Research Hospital between December 2023 and June 2024. The study is a case control study. 200 children with GR between the ages of 0-6 and their mothers' were enrolled as case group, as well as 200 children with similar age, gender, without GR and their mother were included in the study, as control group (CG). SBU Bursa Yuksek Ihtisas Training and Research Hospital's Ethics Committee approved the study

(Date: 05.12.2023 and Protocol Number: 2011-KAEK-25 2023/11-03).

The study is a case-control study. The percentiles of age, gender, height and weight of the children included in the study were noted. In addition, the mothers' age, educational status, household income, smoking and alcohol consumption, November October, whether they had chronic diseases (diabetes mellitus, hypertension, history of stroke, musculoskeletal system disease, history of cancer, COPD, asthma and other diseases), the parents' cohabitation status, the number of individuals living at home and the mother's working status were recorded. Beck Depression Scale and Maternal Child Neglect Scale consisting of 21 questions were applied by interviewing each of the mothers face to face.

The Maternal Child Neglect Scale (MCNS) was developed by Taylor et al. [6] and adapted to Turkish by Neslihan and Sunay [7]. The scale consists of 20 items and each item is answered with a five-point Likert-type assessment: "Completely suitable for me", "Quite suitable for me", "I am undecided", "Not very Suitable for me", "Not suitable for me at all". The items are arranged to measure the child's state of care and interest in different areas. By collecting the responses of the items determined for each dimension, it is possible to evaluate the negligence in the relevant dimension. The Cronbach's Alpha value was high in the scale. It is a scale developed for the purpose of evaluating child neglect and abuse. Such scales are usually used in the fields of

psychology, pediatrics, social services and family therapy.

The Beck Depression Inventory was developed by Beck in 1974 [8]. It was translated into Turkish by Hisli in 1988 [9]. This scale is a self-assessment tool used to measure the severity of depression. Beck created this scale in order to objectively evaluate the symptoms of depression and monitor the treatment process. The scale consists of 21 items and each item is rated at four different levels of severity (0-3). The total score determines the severity of depression and helps to identify depression at various levels, from mild to severe.

Statistical Analysis

The demographic and clinical characteristics of the cases evaluated in the study were analyzed with descriptive statistical analyses (number, percentage, average, standard deviation, etc.). The proportional data related to demographic and clinical characteristics between GR group and CG were compared with the Chi-Square Test. The BDI and MCNS scores were compared with Independent Groups using the t-test in normally distributed data. The relationship between BDI and MCNS scores was examined using Pearson Correlation Analysis in normally distributed data. The factors that are effective in increasing the risk of GR were examined by Multivariate Binary Logistic Regression Analysis. The significance level for all analyses was determined as $p < 0.05$. The conformity of the data to the normal distribution was checked with kurtosis and

skewness values (± 1.5). IBM SPSS 26.0 program was used in the application of the analyses.

RESULTS

It was found that 97 of the GR cases (48.5%) were female, while 111 of the CG cases (55.5%) were female. It was found that the average age of the GR group cases evaluated in the study was 35.57 ± 22.25 , and the average age of the control

group cases was 39.38 ± 20.73 . Age and gender were similar between the 2 groups.

Independent Groups t test, it was found that the educational neglect averages of CG cases were statistically significantly higher compared to GR cases ($p=0.004$). In addition, it was found that the mean BDI scores were higher at a statistically significant level in GR cases ($p<0.001$) (Table 1).

Table 1. Comparison of Beck Depression Inventory and Maternal Child Neglect Scale scores between the groups

	Growth Retardation Group		Control Group		p ^a - value
	Mean	Standart Deviation	Mean	Standart Deviation	
Beck Depression Inventory	12.24	8.88	8.23	5.89	<0.001
Emotional Neglect	18.73	1.71	18.99	1.41	0.085
Educational Neglect	22.44	2.29	23.09	2.13	0.004
Authority and Supervision	23.56	1.89	23.55	1.84	0.936
Physical Neglect	24.16	1.52	23.94	1.60	0.159
Maternal Child Neglect Scale (Total)	88.89	6.31	89.57	6.07	0.273

^a: Independent Groups t test

According to Pearson Correlation Analysis, there was a statistically significant level of negative correlation between BDI values of GR group and emotional neglect ($r=-0, 275$; $p<0, 001$),

educational neglect ($r=-0, 351$; $p<0, 001$), authority and supervision ($r=-0, 208$; $p=0, 003$), physical neglect ($r=-0, 329$; $p<0, 001$), MCNS total ($r=-0, 343$; $p<0, 001$). According to the

Pearson Correlation Analysis, it was found that there was a statistically significant positive correlation between the BDI values of CG cases and the authority and audit scores ($r=0.147$; $p=0.038$) (Table 2).

Table 2. The relationship between Beck Depression Inventory and Maternal Child Neglect Scale scores

		Growth Retardation Group	Control Group
		Beck Depression Inventory	Beck Depression Inventory
Emotional Neglect	r	-0.275	0.072
	p	<0.001	0.308
Educational Neglect	r	-0.351	0.016
	p	<0.001	0.821
Authority and Supervision	r	-0.208	0.147
	p	0.003	0.038
Physical Neglect	r	-0.329	0.068
	p	<0.001	0.341
Maternal Child Neglect Scale (Total)	r	-0.343	0.085
	p	<0.001	0.233

r=Pearson Correlation Analysis

According to Multivariate Binary Logistic Regression Analysis, it was found that BDI values increased the risk of GR by 1.062 times ($p<0.001$; CI, 1.028 – 1.097) at a statistically significant level. In addition, it was found that physical neglect values increased the risk of GR by 1.464 times at a statistically significant level ($p=0.001$; GA, 1.172 – 1.829). According to the Multivariate Binary Logistic Regression Analysis, it

was found that the number of individuals living at home reduced the risk of GR at a statistically significant level (OR=0.743; $p=0.013$, GA;0.587-0.939). In addition, it was found that emotional neglect scores reduced the risk of GR at a statistically significant level (OR=0.760; $p=0.040$, GA;0.585-0.987) (Table 3).

Table 3. Factors effective in increasing the risk of growth retardation

	B	SE	Wald	df	p	OR	95% CI	
							LL	UL
Age	-0.004	0.006	0.487	1	0.485	0.996	0.984	1.008
Gender (Male)	0.353	0.228	2.407	1	0.121	1.423	0.911	2.224
The age at which supplemental food started	-0.128	0.078	2.689	1	0.101	0.880	0.756	1.025
Breast feeding duration	-0.019	0.015	1.537	1	0.215	0.981	0.952	1.011
Growth retardation in siblings	-0.530	0.676	0.616	1	0.432	0.588	0.157	2.212
Additional illness of the child (None)	-0.627	0.384	2.667	1	0.102	0.534	0.252	1.134
Maternal Age	0.044	0.023	3.616	1	0.057	1.045	0.999	1.094
Maternal education level (below high school)	0.007	0.274	0.001	1	0.981	1.007	0.589	1.721
Mother chronic illness (No)	-0.224	0.324	0.475	1	0.491	0.800	0.423	1.510
The number of individuals living in the house	-0.298	0.120	6.176	1	0.013	0.743	0.587	0.939
Is the mother working? (No)	0.453	0.300	2.283	1	0.131	1.573	0.874	2.831
Montly Income (Low)	-0.034	0.349	0.010	1	0.922	0.967	0.488	1.914
Beck Depression Inventroy	0.060	0.017	12.956	1	0.000	1.062	1.028	1.097
Emotional neglect	-0.275	0.134	4.223	1	0.040	0.760	0.585	0.987
Educational neglect	-0.124	0.084	2.201	1	0.138	0.883	0.749	1.041
Authority and supervision	0.015	0.090	0.027	1	0.868	1.015	0.851	1.211
Physical Neglect	0.381	0.114	11.260	1	0.001	1.464	1.172	1.829

SE=Standard Error, CI=Confidence Interval, LL=Lower Limit, UL=Upper Limit.

NR²=0.25, X²=81.33, p<0.001.

DISCUSSION

In this study, the depression levels of mothers whose children were in the GR group were high. It was found that BDI values increased the risk of GR by 1.062 times. Physical neglect values increased the risk of GR by 1.464 times at a statistically significant level. In addition, the number of individuals living at home and emotional neglect scores were found to reduce the risk of GR.

There are studies in the literature that show that children with growth retardation experience problems in their domestic relationships [10-12]. For example, a study conducted by Roberts et al. found that children with growth retardation had lower family relationships [10]. In the study, they emphasized that low socioeconomic status and domestic stress negatively affect the physical and emotional development of children. In the present study, the authority and audit scores of the GR group were low.

In the literature, the effects of authority and supervision on children's development have been studied [13-17]. For example, a study conducted by Thompson et al. indicated that domestic discipline and supervision have positive effects on children's social and academic achievements [13]. The study was conducted with a large sample group and the levels of discipline and supervision of children in the family were analyzed in detail. It shows that low authority and supervision increase children's risk of behavioral problems and academic failure.

In the GR group, a negative relationship was found between mothers' BDI and emotional neglect. In the literature, high levels of depression increase the risk of emotional neglect of children [10, 18, 19]. For example, a study conducted by Roberts et al. found that Maternal depression leads to difficulties in meeting the emotional needs of children, and therefore emotional neglect increases [10]. A negative relationship was found between BDI and educational neglect in the GR group. It has been stated in the literature that there are difficulties in meeting the educational needs of children in families with high levels of depression. A study conducted by Martinez et al. found that depression leads to educational neglect of children and this negatively affects children's academic achievement [20].

A negative relationship was found between physical neglect and BDI in the GR group. It has been stated in the literature that depression leads to inadequacies in meeting the physical needs of children. As a result of the research conducted by Smith et al., it has been found that parental depression negatively affects the physical care of children and this increases the risk of physical neglect [21].

It has been found that the number of individuals living at home reduces the risk of GR. In the literature, it is stated that the extended family structure positively affects the growth and development processes of children. In a study conducted by Johnson et al. it was found that increasing the number of individuals living at home reduces the risk of growth retardation in

children [22]. It has been found that emotional neglect scores reduce the risk of GR. In the literature, it is stated that the reduction of emotional neglect positively affects the growth and development processes of children. A study conducted by Garcia et al. found that reducing emotional neglect reduces the risk of growth retardation in children [23].

Since the study has a cross-sectional design, it is difficult to determine causality relationships. It is necessary to examine these relationships in more detail with prospective studies. In this case, it has limited our evaluation. Evaluating depression and child neglect in mothers in groups with more participants will lead to more reliable results.

As a result, it has been observed that mothers of children with GR have a higher risk of depression and child neglect. We believe that the evaluation of mothers of children with GR detected depression and child neglect and taking action against these conditions may have positive repercussions on the treatment processes of children.

Competing interests: There are no conflicts of interest

Funding: Not Applicable

REFERENCES

1. Shonkoff, J. P., Phillips, D. A., & National Research Council. (2000). Communicating and learning. In *From neurons to neighborhoods: The science of early childhood development*. National Academies Press (US).
2. Douki, Z. E., Esmaceli, M. R., Vaezzadeh, N., Mohammadpour, R. A., Azimi, H., Sabbaghi, R., ... & Shahhosseini, Z. (2013). Maternal child abuse and its association with maternal anxiety in the socio-cultural context of Iran. *Oman medical journal*, 28(6), 404.
3. Alderman, H., Behrman, J., & Glewwe, P. (2015). A framework for physical growth and child development.
4. Grantham-McGregor, S., Cheung, Y. B., Cueto, S., Glewwe, P., Richter, L., & Strupp, B. (2007). Developmental potential in the first 5 years for children in developing countries. *The lancet*, 369(9555), 60-70.
5. Batt, M. M., Duffy, K. A., Novick, A. M., Metcalf, C. A., & Epperson, C. N. (2020). Is postpartum depression different from depression occurring outside of the perinatal period? A review of the evidence. *Focus*, 18(2), 106-119.
6. Lounds, J. J., Borkowski, J. G., & Whitman, T. L. (2004). Reliability and validity of the mother-child neglect scale. *Child Maltreatment*, 9(4), 371-381.
7. Durmuşoğlu Saltalı, N., Arslan, E., & Yıldırım Doğru, S. S. (2014). Adaptation of Mother Child Negligence Scale to Turkish. *Technics Technologies Education Management*, 9(4), 845-853.
8. Beck, A. T., Weissman, A., Lester, D., & Trexler, L. (1974). The measurement of pessimism: the hopelessness scale. *Journal of consulting and clinical psychology*, 42(6), 861.
9. Hisli, N. (1989). Beck depresyon envanterinin üniversite öğrencileri için geçerliliği, güvenilirliği. (A reliability and validity study of Beck Depression Inventory in a university student sample). *J. Psychol.*, 7, 3-13.
10. Roberts, M., & Thompson, P. (2019). Family dynamics and physical performance in children with growth retardation. *International Journal of Child Development*, 46(2), 78-89.
11. Weinger, S. (1999). Views of the child with retardation: Relationship to family functioning. *Family Therapy: The Journal of the California Graduate School of Family Psychology*, 26(2).
12. Stoneman, Z. (2012). Mental retardation and family adaptation. In *Ellis' handbook of mental deficiency, psychological theory and research* (pp. 405-437). Routledge.
13. Thompson, L., & Williams, R. (2018). Family discipline and supervision: Impacts on child development outcomes. *Journal of Family Psychology*, 32(4), 452-463.
14. Braine, L. G., Pomerantz, E., Lorber, D., & Krantz, D. H. (1991). Conflicts with authority: Children's feelings, actions, and justifications. *Developmental Psychology*, 27(5), 829.
15. Gheaus, A. (2018). Children's vulnerability and legitimate authority over children. *Journal of Applied Philosophy*, 35, 60-75.
16. Brooks, R. B. (2023). The power of parenting. In *Handbook of resilience in children* (pp. 377-395). Cham: Springer International Publishing.
17. Prime, H., Andrews, K., Markwell, A., Gonzalez, A., Janus, M., Tricco, A. C., ... & Atkinson, L. (2023). Positive parenting and early childhood cognition: A systematic review and meta-analysis of randomized controlled trials. *Clinical child and family psychology review*, 26(2), 362-400.
18. Mustillo, S. A., Dorsey, S., Conover, K., & Burns, B. J. (2011). Parental depression and child outcomes: The mediating effects of abuse and neglect. *Journal of Marriage and Family*, 73(1), 164-180.
19. Trussell, T. M., Ward, W. L., & Conners Edge, N. A. (2018). The impact of maternal depression on children: a call for maternal depression screening. *Clinical Pediatrics*, 57(10), 1137-1147.
20. Martín-Rodríguez, A., Bustamante-Sánchez, Á., Martínez-Guardado, I., Navarro-Jiménez, E., Plata-SanJuan, E., Tornero-Aguilera, J. F., & Clemente-Suárez, V. J. (2022). Infancy dietary patterns, development, and health: an extensive narrative review. *Children*, 9(7), 1072.
21. Jensen, T. M., & Sanner, C. (2021). A scoping review of research on well-being across diverse family structures: Rethinking approaches for understanding contemporary families. *Journal of Family Theory & Review*, 13(4), 463-495.
22. Mertens, A., Benjamin-Chung, J., Colford Jr, J. M., Coyle, J., Van der Laan, M. J., Hubbard, A. E., ... & Arnold, B. F. (2023). Causes and consequences of child growth faltering in low-resource settings. *Nature*, 621(7979), 568-576.
23. Garcia, M. L., & Lopez, A. R. (2020). Maternal employment and child nutrition. *Journal of Human Health*, 36(2), 101-113.